



Robert S. Williams, MD
Kimberly Petrila, PA-C

Spine Surgery • Orthopedic Surgery • Joint Replacement Surgery • Orthopedic Trauma

Post-Op Total Hip Replacement Care

If you experience nausea or vomiting due to certain anesthetics or pain medications, please contact our office for assistance at home.

Upon discharge, an e-prescription for pain medication will be sent to your designated pharmacy. Use this medication as needed.

You will take aspirin twice daily for one month following your surgery for DVT prophylaxis unless you are already on a blood thinner prescribed by your cardiologist, which you will resume.

Your surgical incision is covered with Prineo, a specialized wound dressing resembling tape. Keep the Prineo in place until your follow-up appointment; no additional wound care is necessary. It's water-resistant so you can shower, but avoid submerging it, as in a bath.

Attending outpatient PT is crucial for the best recovery outcome. Typically, you'll begin therapy within the first two weeks post-op, depending on your insurance. Follow the at-home exercises recommended by the physical therapist from the hospital.

Contact your physician if you experience:

- Temperature over 101.5 degrees or severe chills.
- Severe pain that isn't alleviated by prescribed medications.
- Paleness or coldness of operative extremity.

For symptomatic relief, apply ice to the surgical site for 20 minutes every hour, as needed.

Expect a call from our staff on Monday to check your progress and address any questions you may have.

You will have a **20 lbs weight-bearing restriction (toe touch)** on your operative side for six weeks. You will be discharged with a walker if you do not have one.

Lifelong adherence to total hip precautions is necessary.

Additional instructions: When taking pain medication, ensure you drink plenty of water and take fiber to prevent constipation.

Keep your post-operative appointment at Coastal Orthopedics.

Questions/concerns? Call our office staff (361) 994-1166.